

## S21BAF – NAG Fortran Library Routine Document

**Note.** Before using this routine, please read the Users' Note for your implementation to check the interpretation of bold italicised terms and other implementation-dependent details.

### 1 Purpose

S21BAF returns a value of an elementary integral, which occurs as a degenerate case of an elliptic integral of the first kind, via the routine name.

### 2 Specification

```

real FUNCTION S21BAF(X, Y, IFAIL)
  INTEGER                IFAIL
  real                   X, Y

```

### 3 Description

This routine calculates an approximate value for the integral

$$R_C(x, y) = \frac{1}{2} \int_0^\infty \frac{dt}{\sqrt{t+x(t+y)}}$$

where  $x \geq 0$  and  $y \neq 0$ .

This function, which is related to the logarithm or inverse hyperbolic functions for  $y < x$  and to inverse circular functions if  $x < y$ , arises as a degenerate form of the elliptic integral of the first kind. If  $y < 0$ , the result computed is the Cauchy principal value of the integral.

The basic algorithm, which is due to Carlson [2] and [3], is to reduce the arguments recursively towards their mean by the system:

$$\begin{aligned}
 x_0 &= x, y_0 = y \\
 \mu_n &= (x_n + 2y_n)/3, S_n = (y_n - x_n)/3\mu_n \\
 \lambda_n &= y_n + 2\sqrt{x_n y_n} \\
 x_{n+1} &= (x_n + \lambda_n)/4, y_{n+1} = (y_n + \lambda_n)/4.
 \end{aligned}$$

The quantity  $|S_n|$  for  $n = 0, 1, 2, 3, \dots$  decreases with increasing  $n$ , eventually  $|S_n| \sim 1/4^n$ . For small enough  $S_n$  the required function value can be approximated by the first few terms of the Taylor series about the mean. That is

$$R_C(x, y) = \left( 1 + \frac{3S_n^2}{10} + \frac{S_n^3}{7} + \frac{3S_n^4}{8} + \frac{9S_n^5}{22} \right) / \sqrt{\mu_n}.$$

The truncation error involved in using this approximation is bounded by  $16|S_n|^6/(1 - 2|S_n|)$  and the recursive process is stopped when  $S_n$  is small enough for this truncation error to be negligible compared to the *machine precision*.

Within the domain of definition, the function value is itself representable for all representable values of its arguments. However, for values of the arguments near the extremes the above algorithm must be modified so as to avoid causing underflows or overflows in intermediate steps. In extreme regions arguments are pre-scaled away from the extremes and compensating scaling of the result is done before returning to the calling program.

### 4 References

- [1] Abramowitz M and Stegun I A (1972) *Handbook of Mathematical Functions* Dover Publications (3rd Edition)

- [2] Carlson B C (1978) Computing elliptic integrals by duplication *Preprint* Department of Physics, Iowa State University
- [3] Carlson B C (1988) A table of elliptic integrals of the third kind *Math. Comput.* **51** 267–280

## 5 Parameters

- 1: X — *real* *Input*
- 2: Y — *real* *Input*

*On entry:* the arguments  $x$  and  $y$  of the function, respectively.

*Constraint:*  $X \geq 0.0$  and  $Y \neq 0.0$ .

- 3: IFAIL — INTEGER *Input/Output*

*On entry:* IFAIL must be set to 0,  $-1$  or 1. For users not familiar with this parameter (described in Chapter P01) the recommended value is 0.

*On exit:* IFAIL = 0 unless the routine detects an error (see Section 6).

## 6 Error Indicators and Warnings

If on entry IFAIL = 0 or  $-1$ , explanatory error messages are output on the current error message unit (as defined by X04AAF).

Errors detected by the routine:

IFAIL = 1

On entry,  $X < 0.0$ ; the function is undefined.

IFAIL = 2

On entry,  $Y = 0.0$ ; the function is undefined. On soft failure the routine returns zero.

## 7 Accuracy

In principle the routine is capable of producing full *machine precision*. However round-off errors in internal arithmetic will result in slight loss of accuracy. This loss should never be excessive as the algorithm does not involve any significant amplification of round-off error. It is reasonable to assume that the result is accurate to within a small multiple of the *machine precision*.

## 8 Further Comments

Users should consult the Chapter Introduction which shows the relationship of this function to the classical definitions of the elliptic integrals.

## 9 Example

This example program simply generates a small set of non-extreme arguments which are used with the routine to produce the table of low accuracy results.

## 9.1 Program Text

**Note.** The listing of the example program presented below uses bold italicised terms to denote precision-dependent details. Please read the Users' Note for your implementation to check the interpretation of these terms. As explained in the Essential Introduction to this manual, the results produced may not be identical for all implementations.

```

*      S21BAF Example Program Text
*      Mark 14 Revised.  NAG Copyright 1989.
*      .. Parameters ..
      INTEGER          NOUT
      PARAMETER        (NOUT=6)
*      .. Local Scalars ..
      real             RC, X, Y
      INTEGER          IFAIL, IX
*      .. External Functions ..
      real             S21BAF
      EXTERNAL         S21BAF
*      .. Executable Statements ..
      WRITE (NOUT,*) 'S21BAF Example Program Results'
      WRITE (NOUT,*)
      WRITE (NOUT,*) '      X      Y      S21BAF  IFAIL'
      WRITE (NOUT,*)
      DO 20 IX = 1, 3
         X = IX*0.5e0
         Y = 1.0e0
         IFAIL = 1

*
         RC = S21BAF(X,Y,IFAIL)

*
         WRITE (NOUT,99999) X, Y, RC, IFAIL
20  CONTINUE
      STOP
*
      99999 FORMAT (1X,2F7.2,F12.4,I5)
      END

```

## 9.2 Program Data

None.

## 9.3 Program Results

S21BAF Example Program Results

X	Y	S21BAF	IFAIL
0.50	1.00	1.1107	0
1.00	1.00	1.0000	0
1.50	1.00	0.9312	0

---